



Treatment Tracker

What is LUNSUMIO or LUNSUMIO VELO?

LUNSUMIO (mosunetuzumab-axgb) or LUNSUMIO VELO is a prescription medicine used to treat adults with follicular lymphoma whose cancer has come back or did not respond to previous treatment, and who have already received two or more treatments. It is not known if LUNSUMIO or LUNSUMIO VELO is safe and effective in children.

The conditional approval for this use is based on response rate. There are ongoing studies to establish how well the drug works.

What is the most important information I should know about LUNSUMIO or LUNSUMIO VELO?

LUNSUMIO or LUNSUMIO VELO can cause Cytokine Release Syndrome (CRS), a serious side effect that is common during treatment with LUNSUMIO or LUNSUMIO VELO, and can also be severe or life-threatening.

Get medical help right away if you develop any signs or symptoms of CRS at any time, including:

- fever of 100.4°F (38°C) or higher
- chills
- low blood pressure
- fast or irregular heartbeat
- tiredness or weakness
- difficulty breathing
- headache
- confusion
- feeling anxious
- dizziness or light-headedness
- nausea
- vomiting

Due to the risk of CRS, you will receive LUNSUMIO or LUNSUMIO VELO on a “step-up dosing schedule.”

- The step-up dosing schedule is when you receive smaller “step-up” doses before receiving higher doses of LUNSUMIO or LUNSUMIO VELO during your first cycle of treatment
- If your dose of LUNSUMIO or LUNSUMIO VELO is delayed for any reason, you may need to repeat the “step-up dosing schedule”
- You may receive medicines to help reduce your risk of CRS before your dose

Your healthcare provider will check you for CRS during treatment with LUNSUMIO or LUNSUMIO VELO and may treat you in a hospital if you develop signs and symptoms of CRS. Your healthcare provider may temporarily stop or completely stop your treatment with LUNSUMIO or LUNSUMIO VELO, if you have severe side effects.

What are the possible side effects of LUNSUMIO or LUNSUMIO VELO?

LUNSUMIO or LUNSUMIO VELO can cause serious side effects, including:

- **Neurologic problems. LUNSUMIO or LUNSUMIO VELO can cause serious and life-threatening neurologic problems.** Your healthcare provider will check you for neurologic problems during treatment with LUNSUMIO or LUNSUMIO VELO. Your healthcare provider may also refer you to a healthcare provider who specializes in neurologic problems. Tell your healthcare provider right away if you develop any signs or symptoms of neurologic problems during or after treatment with LUNSUMIO or LUNSUMIO VELO, including:
 - headache
 - numbness and tingling of the arms, legs, hands, or feet
 - dizziness
 - confusion and disorientation
 - difficulty paying attention or understanding things
 - forgetting things or forgetting who or where you are
 - trouble speaking, reading, or writing
 - sleepiness or trouble sleeping
 - tremors
 - loss of consciousness
 - seizures
 - muscle problems or muscle weakness
 - loss of balance or trouble walking
 - tiredness

Please see Important Safety Information, including **Serious Side Effects**, as well as the LUNSUMIO® full [Prescribing Information](#) and [Medication Guide](#) and LUNSUMIO VELO™ full [Prescribing Information](#) and [Medication Guide](#).

Treatment Tracker

Lunsumio
mosunetuzumab-axgb
injection for intravenous use 1 mg | 30 mg

Lunsumio **VELO**
mosunetuzumab-axgb
subcutaneous injection 5 mg | 45 mg

Important dates and details for your treatment journey

Your treatment schedule

LUNSUMIO VELO™ is given as a subcutaneous injection (injected just beneath the skin), and **LUNSUMIO® is given as an intravenous (IV) infusion** (delivered through a needle placed in a vein). Both formulations are designed to be completed in about 6 months (total 8 cycles) or about 1 year (total 17 cycles) depending on your response to treatment. Each cycle is 21 days. The way you start treatment (either as an injection or infusion) is the way it will continue for the rest of your treatment journey.

Treatment may be stopped if it cannot be tolerated or if your disease worsens. If you have not achieved complete remission during the first 6 months of treatment, your healthcare provider may recommend treatment for 6 more months.

In Cycle 1

You'll receive an injection or infusion **3 times** in Cycle 1 (once every 7 days). Each injection will take about 1 minute to receive.* Each infusion will take at least 4 hours.

In Cycles 2-8 or 2-17

You'll receive an injection or infusion on **Day 1** of each following cycle or **once every 21 days**. Each injection will take about 1 minute.* Each infusion will take about 2 hours, if previous infusions in Cycle 1 went well.

Due to the risk of CRS, you will receive LUNSUMIO or LUNSUMIO VELO on a "step-up dosing schedule."

- The step-up dosing schedule is when you receive smaller "step-up" doses before receiving higher doses of LUNSUMIO or LUNSUMIO VELO during your first cycle of treatment
- If your dose of LUNSUMIO or LUNSUMIO VELO is delayed for any reason, you may need to repeat the "step-up dosing schedule"
- You may receive medicines to help reduce your risk of CRS before your dose

*Refers to the injection time and does not include other aspects of treatment; actual clinical time may vary.



Learn more about your treatment schedule

Preparing for your appointment

- You can receive treatment at a local outpatient clinic or infusion center. Some people may be required to go to the hospital to manage certain side effects
- Drink plenty of fluids as it is important to be hydrated before receiving treatment
- Your healthcare provider may have you take medicines before your treatment to reduce possible side effects. These are called premedications and include acetaminophen, antihistamines, and steroid medications



Learn how to prepare for your treatment

After your treatment

Monitor for side effects

LUNSUMIO or LUNSUMIO VELO can cause Cytokine Release Syndrome (CRS), a serious side effect that is common during treatment with LUNSUMIO or LUNSUMIO VELO, and can also be severe or life-threatening.

Get medical help right away if you develop any signs or symptoms of CRS at any time, including:

- | | | |
|-------------------------------------|-------------------------|---------------------------------|
| • fever of 100.4°F (38°C) or higher | • tiredness or weakness | • feeling anxious |
| • chills | • difficulty breathing | • dizziness or light-headedness |
| • low blood pressure | • headache | • nausea |
| • fast or irregular heartbeat | • confusion | • vomiting |

Watch for signs and symptoms of side effects listed throughout. Side effects may be serious, so let your healthcare provider know of any signs or symptoms you may experience.



Find out more about how you might feel after your treatment

Please see Important Safety Information, including **Serious Side Effects**, as well as the LUNSUMIO full [Prescribing Information](#) and [Medication Guide](#) and LUNSUMIO VELO full [Prescribing Information](#) and [Medication Guide](#).

Your LUNSUMIO VELO™ subcutaneous injection or LUNSUMIO® IV infusion schedule
(each cycle is 21 days)

Fill out the section below with your healthcare provider.

Cycle 1	Date	Arrive by:	Location/notes
Cycle 1, Day 1			
Cycle 1, Day 8			
Cycle 1, Day 15			

Cycles 2-8			
Cycle 2			
Cycle 3			
Cycle 4			
Cycle 5			
Cycle 6			
Cycle 7			
Cycle 8			

Cycles 9-17	Cycles 9-17 are for people who need additional treatment, depending on their treatment response.		
Cycle 9			
Cycle 10			
Cycle 11			
Cycle 12			
Cycle 13			
Cycle 14			
Cycle 15			
Cycle 16			
Cycle 17			

Please see Important Safety Information, including **Serious Side Effects**, as well as the LUNSUMIO full [Prescribing Information](#) and [Medication Guide](#) and LUNSUMIO VELO full [Prescribing Information](#) and [Medication Guide](#).



The goal of this resource is to help keep track of side effects after injections or infusions. Your healthcare provider may fill in the top section. Then, you can fill in the dates, times, and measurements. The chart below may not include all of the information you may need. On the next page, you'll find additional space for you and your healthcare provider to fill in any other observations you record after your injections or infusion.

Get medical help right away if you develop any signs or symptoms of CRS at any time.

Monitoring notes:

Measure (eg, temperature): _____	Measure: _____	Measure: _____
Frequency: _____	Frequency: _____	Frequency: _____
Call healthcare provider if: _____	Call healthcare provider if: _____	Call healthcare provider if: _____
_____	_____	_____
Go to emergency department if: _____	Go to emergency department if: _____	Go to emergency department if: _____
_____	_____	_____

[illegible]

Download the Wallet Card and carry it with you to help your healthcare providers understand what treatments you are taking.

Lunsumio
mosunetuzumab-axgb
injection for intravenous use 1 mg | 30 mg

Lunsumic  **VELO**[™]
mosunetuzumab-axgb
subcutaneous injection 5 mg | 45 mg

Additional monitoring notes

Use this space to record any side effects, symptoms, changes, or important notes to share about your treatment experience.

Important contact information

Infusion center contact information

Name of healthcare provider: _____

Contact during business hours: _____

Contact after business hours: _____

Emergency department contact information

Name: _____

Address: _____

Phone: _____



For information on financial assistance,
visit LUNSUMIO.com/support

Important Safety Information (cont'd)

What are the possible side effects of LUNSUMIO or LUNSUMIO VELO? (cont'd)

- **Serious infections.** LUNSUMIO or LUNSUMIO VELO can cause serious infections that may lead to death. Your healthcare provider will check you for signs and symptoms of infection before and during treatment. Tell your healthcare provider right away if you develop any signs or symptoms of infection during treatment with LUNSUMIO or LUNSUMIO VELO, including:
 - fever of 100.4°F (38°C) or higher
 - cough
 - chest pain
 - tiredness
 - shortness of breath
 - painful rash
 - sore throat
 - pain during urination
 - feeling weak or generally unwell

Please see Important Safety Information, including **Serious Side Effects**, as well as the LUNSUMIO® full [Prescribing Information](#) and [Medication Guide](#) and LUNSUMIO VELO™ full [Prescribing Information](#) and [Medication Guide](#).

Lunsumio[®]
mosunetuzumab-axgb
injection for intravenous use 1 mg | 30 mg

Lunsumio[®] **VELO**[™]
mosunetuzumab-axgb
subcutaneous injection 5 mg | 45 mg

Important Safety Information (cont'd)

What are the possible side effects of LUNSUMIO or LUNSUMIO VELO? (cont'd)

- **Hemophagocytic lymphohistiocytosis (HLH).** LUNSUMIO or LUNSUMIO VELO can cause overactivity of the immune system, a condition called hemophagocytic lymphohistiocytosis. HLH can be life-threatening and has led to death in people treated with LUNSUMIO or LUNSUMIO VELO. Your healthcare provider will check you for HLH especially if your CRS lasts longer than expected. Signs and symptoms of HLH include:
 - fever
 - enlarged spleen
 - easy bruising
 - low blood cell counts
 - liver problems
- **Low blood cell counts.** Low blood cell counts are common during treatment with LUNSUMIO or LUNSUMIO VELO and can also be serious or severe. Your healthcare provider will check your blood cell counts during treatment with LUNSUMIO or LUNSUMIO VELO. LUNSUMIO or LUNSUMIO VELO can cause the following low blood cell counts:
 - **low white blood cell counts (lymphopenia [for LUNSUMIO VELO only] and neutropenia).** Low white blood cells can increase your risk for infection
 - **low red blood cell counts (anemia).** Low red blood cells can cause tiredness and shortness of breath
 - **low platelet counts (thrombocytopenia).** Low platelet counts can cause bruising or bleeding problems
- **Growth in your tumor or worsening of tumor-related problems (tumor flare).** LUNSUMIO or LUNSUMIO VELO can cause serious or severe worsening of your tumor. Tell your healthcare provider if you develop any of these signs or symptoms of tumor flare during your treatment with LUNSUMIO or LUNSUMIO VELO:
 - chest pain
 - cough
 - trouble breathing
 - tender or swollen lymph nodes
 - pain or swelling at the site of the tumor

Your healthcare provider may temporarily stop or permanently stop treatment with LUNSUMIO or LUNSUMIO VELO if you develop severe side effects.

The most common side effects of LUNSUMIO include: CRS, tiredness, rash, headache, fever, muscle pain, cough, itching, and numbness, tingling, or pain in the hands or feet (nerve damage).

The most common side effects of LUNSUMIO VELO include: injection site reactions, tiredness, rash, CRS, COVID-19, muscle and joint pain, and diarrhea.

The most common severe abnormal blood test results with LUNSUMIO include: decreased phosphate, increased glucose, and increased uric acid levels.

The most common severe abnormal blood test results with LUNSUMIO VELO include: decreased white blood cell counts and increased uric acid levels.

Before receiving LUNSUMIO or LUNSUMIO VELO, tell your healthcare provider about all of your medical conditions, including if you:

- have ever had an infusion reaction after receiving LUNSUMIO
- have an infection or have had an infection in the past which lasted a long time or keeps coming back
- have or have had Epstein-Barr Virus
- are pregnant or plan to become pregnant. LUNSUMIO or LUNSUMIO VELO may harm your unborn baby. Tell your healthcare provider right away if you become pregnant or think you may be pregnant during treatment with LUNSUMIO or LUNSUMIO VELO

Females who are able to become pregnant:

- your healthcare provider should do a pregnancy test before you start treatment with LUNSUMIO or LUNSUMIO VELO
- use an effective method of birth control (contraception) during your treatment and for 3 months after the last dose of LUNSUMIO or LUNSUMIO VELO
- are breastfeeding or plan to breastfeed. It is not known if LUNSUMIO or LUNSUMIO VELO passes into your breast milk. Do not breastfeed during treatment and for 3 months after the last dose of LUNSUMIO or LUNSUMIO VELO

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

What should I avoid while receiving LUNSUMIO or LUNSUMIO VELO?

Do not drive, operate heavy machinery, or do other dangerous activities if you develop dizziness, confusion, tremors, sleepiness, or any other symptoms that impair consciousness until your signs and symptoms go away. These may be signs and symptoms of CRS or neurologic problems.

These are not all of the possible side effects of LUNSUMIO or LUNSUMIO VELO. Talk to your healthcare provider for more information about the benefits and risks of LUNSUMIO or LUNSUMIO VELO.

You may report side effects to the FDA at (800) FDA-1088 or www.fda.gov/medwatch. You may also report side effects to Genentech at (888) 835-2555.

Please see Important Safety Information, including **Serious Side Effects**, as well as the LUNSUMIO® full [Prescribing Information](#) and [Medication Guide](#) and LUNSUMIO VELO™ full [Prescribing Information](#) and [Medication Guide](#).

